

AUBURN NY ORAL SURGERY

Sandeep Singla DDS, MD

Rinil Patel DDS

Edward Woodbine DDS

www.auburnnyoralsurgery.com

183 Genesee Street

Auburn, NY 13021

Tel: (315) 253-7384

Fax: (315) 253-7426

Post-Operative Instructions for Sinus Lift or Augmentation Surgery

A certain amount of swelling, pain and bleeding is normal.

The following instructions are provided to make your postoperative recovery period as short and pleasant as possible.

You will be having a Sinus Lift if the area in the upper jaw that you are to have an implant(s) currently has insufficient bone because of an enlarged sinus. This procedure will augment the deficient area in the maxillary sinus. The bone that is being grafted will most commonly be coming from freeze dried bone and artificial synthetic bone. Other sites may be used for donation of your own bone as previously discussed with you by our doctors. Because of this you may have two post-surgical wounds: The donor site and the recipient site.

DO NOT UNDER ANY CIRCUMSTANCES. BLOW YOUR NOSE FOR THE NEXT FOUR (4) WEEKS. This may be longer if indicated. You may sniff all you like but NO BLOWING.

Bite down and maintain firm pressure on the gauze pack that was placed at the end of your surgery. There may be two areas of gauze in your mouth. Do this for at least 30 minutes. At that time remove the pack and replace it with another gauze sponge as provided. Repeat this as necessary. There should not be excessive bleeding. Bright red blood rapidly pooling in the area is not normal. If this occurs and does not decrease with pressure please call the office immediately. There may be a small amount of nasal discharge. It may be clear or lightly bloody. Excessive bleeding from the nose should be reported immediately.

If you were asleep for the surgery be sure to eat a soft nutritious food prior to taking any medications. This will decrease the chances of post-surgical nausea. You were prescribed an antibiotic prior to surgery this should be continued until all the pills have been taken. You were prescribed a decongestant; take this until it is all gone. Take your pain medications as needed in accordance with the directions. Codeine, hydrocodone or Demerol can be supplemented with Advil or ibuprofen type medications. Be sure you have no allergies that the doctor does not know about.

Do not rinse or spit for a minimum of 72 hours and longer if possible. This tends to disturb the blood clot and can prolong bleeding, open the wound and slow healing. There should be no blood in your mouth.

Saliva can be swallowed. Do not brush your teeth next to the surgery site for 48 hours or only brush in areas away from your graft surgery. When expectorating, BE GENTLE. If you used a prescribed mouthwash prior to surgery, it should be continued starting 24 hours after your surgery. Place the mouthwash in the area of the surgery and let it sit 30-60 seconds. Do not rinse with it until after you have returned for suture removal.

Apply ice immediately in the areas where the surgery was done. Since your graft was in the upper jaw you can expect swelling and discomfort up into the nose and even under the eyes and cheeks. Bruising and discoloration is

